Epilepsy Res., 9 (1991) 253–255
Elsevier

Author Index to Volume 9 (1991)

Anderson, W.W., see Churn, S.B., 211
Antoniadis, G., see Kostopoulos, G.L., 184
Antonini, A., see Hajek, M., 44
Avanzini, G., see Isokawa, M., 242

Babb, T.L., see Isokawa, M., 242
Baker, G.A., see Smith, D.F., 231
Blacklaw, J., see Larkin, J.G., 71
Blosser, J.C., see Garske, G.E., 161
Boehrer, A., see Vergnes, M., 97
Breese, G.R., see McCown, T.C., 132
Brinciotti, M., see Matricardi, M., 145
Brodie, M.J., see Larkin, J.G., 71
Bromfield, E.B., Ludlow, C.L., Sedory, S., Leiderman, D.B. and Theodore, W.H., Cerebral activation during speech discrimination in temporal lobe epilepsy, 49

Calvo, J.M., Amygdaloid kindling during wakefulness and paradoxical sleep in the cat. 1. Inhibitory influence of paradoxical sleep on kindling development, 113
Calvo, J.M. and Fernández-Mas, R., Amygdaloid kindling during wakefulness and paradoxical sleep in the cat. 2. Sleep organization changes produced by kindling development, 175
Chadwick, D.W., see Smith, D.F., 231
Chapman, A.G., Smith, S.E. and Meldrum, B.S., The anticonvulsant effect of the non-NMDA antagonists, NBQX and GYKI 52466, in mice, 92
Churn, S.B., Anderson, W.W. and DeLorenzo, R.J., Exposure of hippocampal slices to magnesium-free medium produces epileptiform activity and simultaneously decreases calcium and calmodulin-dependent protein kinase II activity, 211
Coenen, A.M.L., Van Luijtelaar, E.L.J.M., De Bruijn, S.F.T.M., Declerck, A.C., Renier, W.O. and Vossen, J.M.H., Disturbances in time estimation during absence seizures in children, 148
Coenen, A.M.L., see Drinkenburg, W.H.I.M., 218
Coenen, A.M.L., see Ramakers, G.M.J., 127

De Beukelaar, F., see Rentmeester, T., 59, 65
De Bruijn, S.F.T.M., see Coenen, A.M.L., 148
Declerck, A.C., see Coenen, A.M.L., 148
DeLorenzo, R.J., see Churn, S.B., 211
Depaulis, A., see Lannes, B., 79
Depaulis, A., see Vergnes, M., 97
Dewey, M., see Smith, D.F., 231
Dickinson, S.L., see Jackson, H.C., 121
Drinkenburg, W.H.I.M., Coenen, A.M.L., Vossen, J.M.H. and Van Luijtelaar, E.L.J.M., Spike-wave discharges and sleep-wake states in rats with absence epilepsy, 218

Edgar, P.P., see McCown, T.C., 132

Fassbender, C.P., see Löscher, W., 195
Fernández-Mas, R., see Calvo, J.M., 175
Finch, D.M., see Isokawa, M., 242
Freedman, L.R., see Garske, G.E., 161

Garske, G.E., Palmer, G.C., Napier, J.J., Griffith, R.C., Freedman, L.R., Harris, E.W., Ray, R., McCreedy, S.A., Blosser, J.C., Woodhead, J.H., White, H.S. and Swinyard, E.A., Preclinical profile of the anticonvulsant remacemide and its enantiomers in the rat, 161
Gioanni, H., see Gioanni, Y., 19
Gioanni, Y., Gioanni, H. and Mitrovic, N., Seizures can be triggered by stimulating non-cortical structures in the quaking mutant mouse, 19
Griffith, R.C., see Garske, G.E., 161

Hajek, M., Antonini, A., Leenders, K.L. and Wieser, H.-G., Epilepsia partialis continua studied by PET, 44
Hamberger, A., Nyström, B., Larsson, S., Silfvenius, H. and Nordborg, C., Amino acids in the neuronal microenvironment of focal human epileptic lesions, 32
Harris, E.W., see Garske, G.E., 161
Hebebrand, J., see Schmitz, E., 105
Hulsman, J., see Rentmeester, T., 59, 65
Hönack, D., see Löscher, W., 195
Hönack, D., see Schmitz, E., 105

Isojärvi, J.I.T., Pakarinen, A.J. and Myllylä, V.V., A prospective study of serum sex hormones during carbamazepine therapy, 139
Isokawa, M., Avanzini, G., Finch, D.M., Babb, T.L. and Levesque, M.F., Physiologic properties of human dentate granule cells in slices prepared from epileptic patients, 242
Ito, M., see Oguro, K., 225

Jackson, H.C., Ripley, T.L., Dickinson, S.L. and Nutt, D.J., Anticonvulsant activity of the imidazoline, 6,7-benzoidazoxan, 121
Jacoby, A., see Smith, D.F., 231
Janssen, A., see Rentmeester, T., 59, 65

Kempf, E., see Lannes, B., 79
Kostopoulos, G. and Antoniadis, G., A comparison of recurrent inhibition and of paired-pulse facilitation in hippocampal slices from normal and genetically epileptic mice, 184

Lannes, B., Vergnes, M., Marescaux, C., Depaulis, A., Micheletti, G., Warter, J.-M. and Kempf, E., Lesions of nor-

adrenergic neurons in rats with spontaneous generalized non-convulsive epilepsy, 79
Larkin, J.G., McKee, P.J.W., Blacklaw, J., Thompson, G.G., Morgan, I.C. and Brodie, M.J., Nimodipine in refractory epilepsy: a placebo-controlled, add-on study, 71
Larsson, S., see Hamberger, A., 32
Leenders, K.L., see Hajek, M., 44
Leiderman, D.B., see Bromfield, E.B., 49
Levesque, M.F., see Isokawa, M., 242
Lok, S.Y., see Millan, M.H., 86
Löscher, W. and Nolting, B., The role of technical, biological and pharmacological factors in the laboratory evaluation of anticonvulsant drugs. IV. Protective indices, 1
Löscher, W., Hönack, D., Nolting, B. and Fassbender, C.P., Trans-2-en-valproate: reevaluation of its anticonvulsant efficacy in standardized seizure models in mice, rats and dogs, 195
Löscher, W., see Schmitz, E., 105
Lothman, E.W., Williamson, J.M. and VanLandingham, K.E., Intraperitoneal phenytoin suppresses kindled responses: effects on motor and electrographic seizures, 11
Ludlow, C.L., see Bromfield, E.B., 49

Marescaux, C., see Lannes, B., 79
Marescaux, C., see Vergnes, M., 97
Matricardi, M., Brinciotti, M. and Paciello, F., Reading epilepsy with absences, television-induced seizures, and pattern sensitivity, 145
McCown, T.C., Edgar, P.P., Schwartz, R.D. and Breese, G.R., Unilateral kindling of the inferior collicular cortex does not transfer to the contralateral seizure sensitive site or alter [^{3}H]flunitrazepam and [^{35}S]TBPS binding, 132
McCreedy, S.A., see Garske, G.E., 161
McKee, P.J.W., see Larkin, J.G., 71
Meijer, J., see Rentmeester, T., 59, 65
Meldrum, B.S., see Chapman, A.G., 92
Meldrum, B.S., see Millan, M.H., 86
Micheletti, G., see Lannes, B., 79
Mikawa, H., see Oguro, K., 225
Millan, M.H., Obrenovitch, T.P., Sarna, G.S., Lok, S.Y., Symon, L. and Meldrum, B.S., Changes in rat brain extracellular glutamate concentration during seizures induced by systemic picrotoxin of focal bicuculline injection: an in vivo dialysis study with on-line enzymatic detection, 86
Mitrovic, N., see Gioanni, Y., 19
Morgan, I.C., see Larkin, J.G., 71
Mutoh, K., see Oguro, K., 225
Myllylä, V.V., see Isojärvi, J.I.T., 139

Napier, J.J., see Garske, G.E., 161
Nolting, B., see Löscher, W., 1, 195
Nordborg, C., see Hamberger, A., 32
Nutt, D.J., see Jackson, H.C., 121
Nyström, B., see Hamberger, A., 32

Obrenovitch, T.P., see Millan, M.H., 86
Oguro, K., Ito, M., Tsuda, H., Mutoh, K., Shiraishi, H., Shirasaka, Y. and Mikawa, H., Association of NMDA receptor sites and seizures of El mice, 225
Overweg, J., see Rentmeester, T., 59, 65

Paciello, F., see Matricardi, M., 145
Pakarinen, A.J., see Isojärvi, J.I.T., 139
Palmer, G.C., see Garske, G.E., 161
Peeters, B.W.M.M., see Ramakers, G.M.J., 127

Ramakers, G.M.J., Peeters, B.W.M.M., Vossen, J.M.H. and Coenen, A.M.L., CNQX, a new non-NMDA receptor antagonist, reduces spike wave discharges in the WAG/Rij rat model of absence epilepsy, 127
Ray, R., see Garske, G.E., 161
Reichelt, R., see Schmitz, E., 105
Renier, W.O., see Coenen, A.M.L., 148
Rentmeester, T., Janssen, A., Hulsman, J., Scholtes, F., Van der Kleij, B., Overweg, J., Meijer, J. and De Beukelaar, F., A double-blind, placebo-controlled evaluation of the efficacy and safety of loreclezole as add-on therapy in patients with uncontrolled partial seizures, 59
Rentmeester, T., Janssen, A., Hulsman, J., Scholtes, F., Van der Kleij, B., Overweg, J., Meijer, J. and De Beukelaar, F., Long-term evaluation of the efficacy and safety of loreclezole as add-on therapy in patients with uncontrolled partial seizures: a 1-year open follow-up, 65
Ripley, T.L., see Jackson, H.C., 121

Sarna, G.S., see Millan, M.H., 86
Schmitz, E., Löscher, W., Hönack, D., Reichelt, R. and Hebebrand, J., Kindling does not induce persistent changes in fluorographic labeling patterns of benzodiazepine binding proteins in various rat brain regions, 105
Scholtes, F., see Rentmeester, T., 59, 65
Schwartz, R.D., see McCown, T.C., 132
Sedory, S., see Bromfield, E.B., 49
Shiraishi, H., see Oguro, K., 225
Shirasaka, Y., see Oguro, K., 225
Silfvenius, H., see Hamberger, A., 32
Smith, D.F., Baker, G.A., Dewey, M., Jacoby, A. and Chadwick, D.W., Seizure frequency, patient-perceived seizure severity and the psychosocial consequences of intractable epilepsy, 231
Smith, S.E., see Chapman, A.G., 92
Swinyard, E.A., see Garske, G.E., 161
Symon, L., see Millan, M.H., 86

Theodore, W.H., see Bromfield, E.B., 49
Thompson, G.G., see Larkin, J.G., 71
Tsuda, H., see Oguro, K., 225

Van Luijtelaar, E.L.J.M., see Coenen, A.M.L., 148
Van Luijtelaar, E.L.J.M., see Drinkenburg, W.H.I.M., 218
Van der Kleij, B., see Rentmeester, T., 59, 65
VanLandingham, K.E., see Lothman, E.W., 11
Vergnes, M., Marescaux, C., Boehrer, A. and Depaulis, A., Are rats with genetic absence epilepsy behaviorally impaired?, 97
Vergnes, M., see Lannes, B., 79

Vossen, J.M.H., see Coenen, A.M.L., 148
Vossen, J.M.H., see Drinkenburg, W.H.I.M., 218
Vossen, J.M.H., see Ramakers, G.M.J., 127

Warter, J.-M., see Lannes, B., 79
White, H.S., see Garske, G.E., 161
Wieser, H.G., Comparison of valproate concentrations in human plasma, CSF and brain tissue after administration of different formulations of valproate or valpromide, 154
Wieser, H.-G., see Hajek, M., 44
Williamson, J.M., see Lothman, E.W., 11
Woodhead, J.H., see Garske, G.E., 161